

# Properties of $\tau$ lepton

ON BEHALF OF LEP COLLABORATIONS

*Vadym Zhuravlov*

*JINR Dubna*

*and*

*MPI für Physik Munich*

# New results from LEP

## ➤ $\tau$ decay

- ✓  $\tau \rightarrow \text{hadrons}$     DELPHI
- ✓ lifetime            DELPHI
- ✓  $\nu_\tau$  mass limit from  
 $\tau \rightarrow 5\pi$             DELPHI

## ➤ $\tau$ production


- ✓  $\gamma\gamma \rightarrow \tau\tau$     DELPHI and L3
  - cross-section
  - electromagnetic moments
- ✓  $ee \rightarrow \tau\tau$ 
  - weak dipole momentsALEPH

*Results are preliminary!*

# $ee \rightarrow \tau\tau$ selection at LEP

- Two low multiplicity "back-to-back" jets.  
Cut on charged multiplicity rejects  $Z \rightarrow qq$  background
- $\gamma\gamma$  and  $Z \rightarrow ee, Z \rightarrow \mu\mu$  backgrounds rejected by cuts on visible energy
- Tau selection based on event kinematic properties only.  
No particle identification

	ALEPH	DELPHI	L3	OPAL
efficiency	80 %	62 %	62 %	75%
background	2 %	3 %	2 %	3%



# Hadronic Branching Ratios


## Delphi 92 - 95

### ➤ Charged multiplicity:



- photon conversion: 7 % of  $\gamma$ 's converted before TPC
- Dalitz decays: 1 % of  $\pi^0$  decay to  $e^+ e^- \gamma$
- nuclear reinteraction: 3 % of hadrons interact before TPC
- $\delta$ -rays, two-track resolution, ...

### ➤ $\pi^0$ reconstruction:



- type I  
single EM cluster:  
analysis of cluster structure
- type II  
2 EM clusters
- type III  
EM cluster + converted  $\gamma$
- type IV  
EM cluster + EM energy  
associated to track




# Hadronic branching ratios: 1-prong



# Hadronic branching ratios: 3- and 5- prongs



# $\tau$ lifetime: decay vertex reconstruction method



## 1. Decay length determination:

distance between center of beam crossing region  
and decay vertex.  $\sigma = \sigma_{\text{prod}} \oplus \sigma_{\text{decay}}$

## 2. Conversion to decay time:

dividing by  $\gamma \beta c \sin \theta_\tau$

## 3. Fit of $\tau$ lifetime to the data:

$$\log L = \sum \log (f_{\text{vtx}} \otimes f_{\text{res}})$$

$$f_{\text{vtx}} = (1 - b_f - b_h) E(t | \tau) + b_f E(t | 0.75 \tau) + b_h \delta(t)$$


$$f_{\text{res}} = (1 - f_2 - f_3) G(t | k_1 \sigma) + f_2 G(t | k_2 \sigma) + f_3 G(t | k_3 \sigma)$$

$$1 : k_2 : k_3 = 1.0 : 1.6 : 5.1$$

$$1 : f_2 : f_3 = 1 : 0.25 : 0.007$$

$\tau_{\text{3-prong}} = 288.6 \pm 2.4_{\text{(stat)}} \pm 1.4_{\text{(syst)}} \text{ fs}$

**DELPHI 1991 - 1995 data sample**  
15427 3v1 and 2101 3v3  
(total 19629 vertices)



# $\tau$ lifetime: impact parameter difference method

Impact parameter in transverse plane:

$$d = L \sin \theta_\tau \sin(\phi - \phi_\tau)$$

Impact parameter difference:

$$d_+ - d_- = L_+ \sin \theta_{\tau^+} \sin(\phi_+ - \phi_{\tau^+}) - L_- \sin \theta_{\tau^-} \sin(\phi_- - \phi_{\tau^-})$$

Averaging over decay lengths

$$\langle d_+ - d_- \rangle = \langle L \rangle (\phi_+ - \phi_- + \pi) \sin \theta_\tau = \langle L \rangle \Delta \phi \sin \theta_\tau$$

$$\langle L \rangle = \gamma \beta c \tau_\tau$$

Impact parameter difference is proportional to projected acoplanarity with a proportionality constant  $\sim$  lifetime


Lifetime is extracted by straight line fit to

$$Y = d_+ - d_-$$


as a function of

$$X = \Delta \phi \sin \theta_\tau$$

$$\tau_{ipd} = 291.4 \pm 3.6 \text{ (stat)} \pm 1.5 \text{ (syst)} \text{ fs}$$



1994 and 1995  
26036 events



# $\tau$ lifetime: miss distance method

Miss distance

$$d_{\text{miss}} = d_+ + d_-$$

Lifetime is determined by fit to the observed  $d_{\text{miss}}$  distribution with probability density function

$$f = f_{\text{phys}} \otimes f_{\text{res}}$$

Physics function:



$$f_{\text{phys}} = f_+(d_+) \otimes f_-(d_-)$$

Resolution function:

$$f_{\text{res}} = f_1 G(\sigma_1) + f_2 G(\sigma_2) + f_3 G(\sigma_3)$$

$f_1, f_2, f_3$  and  $\sigma_1, \sigma_2, \sigma_3$  are functions of  $p$  and  $\theta$ , calibrated separately for hadronic and leptonic decays

$\tau_{\text{miss}} = 292.0 \pm 2.3 \text{ (stat)} \pm 2.1 \text{ (stat)} \text{ fs}$



# $\tau$ lifetime

Average:

IPD and MD (1991 -1995 ) 36% correlation


$$\tau_{\text{1-prong}} = 292.3 \pm 1.8_{\text{(stat)}} \pm 1.2_{\text{(syst)}} \text{ fs}$$

1-prong and 3-prong

5% correlation

$$\tau_\tau = 290.9 \pm 1.4_{\text{(stat)}} \pm 1.0_{\text{(syst)}} \text{ fs}$$

Most precise measurement to date!



# $\nu_\tau$ mass limit from $\tau \rightarrow 5\pi$

Likelihood function:

$$L = \prod_i f(m_i E_i) \otimes R(m_i E_i) \otimes \varepsilon(m_i E_i)$$

$$m_i = m_{5\pi} \quad E_i = E_{5\pi}$$

Fit region:

$$m > 1.6 \text{ GeV}/c^2$$

$$0.85 < E / E_{\text{beam}} < 1.10$$


Typical resolution:

$$m: 16 \text{ MeV}/c^2$$


$$E: 500 \text{ MeV}$$

Events in fit region: 15 ( 6 % bkg )

Spectral function:




**DELPHI 1992 - 1995**  
**47 5 $\pi$  vertices**  
background - 24 %  
(mainly qq and 5 $\pi\pi^0$ )





# Summary of $\nu$ mass limits


	3 $\pi$		5 $\pi$		Combined
	Events	Limit <sub>95</sub>	Events	Limit <sub>95</sub>	
ALEPH	2939	22.3 MeV	55	21.5 MeV	18.2 MeV
DELPHI	12 K	28 MeV	15	48.0 MeV	
OPAL	2514	35.3 MeV	22	43.2 MeV	27.6 MeV
ARGUS			19	31 MeV	
CLEO	17 K (3 $\pi\pi^0$ )	28 MeV (3 $\pi\pi^0$ )	55	30 MeV	

DELPHI expected limit  $\approx$  30 MeV

$\gamma\gamma \rightarrow \tau\tau$ 


	DELPHI	L3
Data set	183 - 208 GeV	189 - 208 GeV
Mode	$\tau^+ \rightarrow e^+ \nu \bar{\nu}$ $\tau^- \rightarrow \text{non } e^-$	$\tau^+ \rightarrow e^+ \nu \bar{\nu}$ $\tau^- \rightarrow \pi^+ \pi^0 \nu$
e ID	dE/dx	BGO shower
non-e ID	dE/dx	charged track + 2 $\gamma$ $m(\gamma\gamma) = m(\pi^0)$ $m(\pi^0 \pi^\pm) = m(\rho)$
Sel.eff	0.8 %	0.1 %
Selected events	2154	266
background:	10 %	4 %





$$F_1(q^2)\gamma_\mu + iF_2(q^2)\sigma_{\mu\nu}\frac{q_\nu}{2m_\tau} + F_3(q^2)\gamma^5\sigma^{\mu\nu}\frac{q_\nu}{2m_\tau}$$

At  $q^2=0$

F2 - anomalous magnetic moment  $a_\tau$

F3 - dipole moment  $d_\tau$

SM values:  $a_\tau = 1.1773(3) \cdot 10^{-3}$   
 $d_\tau \approx 0$

Limits from  $\gamma\gamma \rightarrow \tau\tau$  cross-sections:

DELPHI (95% C.L.)

$$-0.017 < a_\tau < 0.019$$


$$|d_\tau| < 3.8 \cdot 10^{-16} e \cdot \text{cm}$$

L3 (68% C.L.)

$$-0.062 < a_\tau < 0.044$$

$$|d_\tau| < 6.7 \cdot 10^{-16} e \cdot \text{cm}$$


$\gamma\gamma \rightarrow \tau\tau$ : electromagnetic moments



# Weak dipole moments

Neutral current:  $\bar{\tau} \left[ v_\tau \gamma^\mu - a_\tau \gamma^\mu \gamma^5 + i \frac{\mu_\tau^Z}{2m_\tau} \sigma^{\mu\nu} q_\nu + \frac{d_\tau^Z}{2m_\tau} \gamma^5 \sigma^{\mu\nu} q_\nu \right] \tau$

Charged current:  $\bar{\nu} \left[ \gamma_\mu \frac{1 - \gamma^5}{2} - i \frac{\kappa_\tau^W}{2m_\tau} \sigma^{\mu\nu} q_\nu \frac{1 + \gamma^5}{2} - \frac{d_\tau^W}{2m_\tau} \gamma^5 \sigma^{\mu\nu} q_\nu \frac{1 + \gamma^5}{2} \right] \tau$



# Weak dipole moments

ALEPH measurement:

- 1990 - 1995
- $\tau \rightarrow \pi, \rho, a_1$  decays
- fit to spin-dependent differential cross-section

$$\text{Re}(\mu_\tau) \quad (-0.33 \pm 0.42 \pm 0.26) \cdot 10^{-3}$$

$$\text{Im}(\mu_\tau) \quad (-0.99 \pm 0.80 \pm 0.61) \cdot 10^{-3}$$

$$\text{Re}(d_\tau) \quad (-0.59 \pm 2.14 \pm 1.26) \cdot 10^{-18} e \cdot \text{cm}$$

$$\text{Im}(d_\tau) \quad (-0.45 \pm 4.00 \pm 4.01) \cdot 10^{-18} e \cdot \text{cm}$$

*Most precise measurement to date!*

- $d_\tau^W$  not measured yet ...
- Simultaneous fit to all moments  $\mu_\tau^Z, d_\tau^Z, \kappa_\tau^W, d_\tau^W$  can be done
- Experimental bounds on charged current moments  $\kappa_\tau^W$  and  $d_\tau^W$  are not listed in PDG

# Conclusion



All LEP experimental data are analysed,  
almost all results are published,  
preliminary results are close to completion