A New Generation of CTEQ Parton Distribution Functions with Uncertainty Analysis

Daniel Stump

Michigan State University

CTEQ6

J. Pumplin, D. Stump, J. Huston, H.L. Lai,
P. Nadolsky, W.K. Tung

hep-ph/0201195

Parton Distribution Functions

For any short-distance process, e.g., *ep* DIS,



the factorization theorem of QCD (*schematically*)

$$\sigma(Q) = \int \; f_i(x,Q)\; \hat{\sigma}_i(x,Q)dx$$

relates experimental $\sigma(Q)$ and perturbatively calculated $\hat{\sigma}_i(x,Q)$.

Global analysis

- Use data from many processes to determine the universal PDF's.
- Parametrize the $f_i(x,Q_0)$ at Q_0=1.3 GeV, with ~20 fitting parameters $\{a_1, a_2, \ldots, a_n\}$.

What is new in the CTEQ6 analysis?

- New Data
  - H1 and ZEUS : deep inelastic ep and $\bar{e}p$ scattering
  - DØ : $p\bar{p} \rightarrow jet$ cross section, as a function of η and E_T.

- New methods of analysis
  - For systematic errors … The published systematic errors are included in the fitting procedure.
  - For uncertainties … Methods are available to evaluate the uncertainties of the PDF's and their predictions.

Data used in the CTEQ6 global analysis

Data for which detailed systematic errors have been published and used in the fit….

process	data set	χ^2_e/N_e
DIS μp	BCDMS p	378/339
DIS μd	BCDMS d	280/251
DIS ep	H1a	99/104
DIS ep	H1b	129/126
DIS ep	ZEUS	263/229
DIS μp	NMC F2p	305/201
DIS μd	NMC F2d/p	112/123
$p\bar{p} \to$ jet	DØ jet	69/90
$p\bar{p} \to$ jet	CDF jet	49/33

Other data used in the global analysis but without systematics….

process	data set	χ^2_e/N_e
DIS ν Fe	CCFR	150/156
DY pp	E605	95/119
DY pd/pp	E866	6/15
$p\bar{p} \to W$	CDF W	10/11

Overall $\chi^2/N = 1954/1811$

χ^2 minimization

The simplest fitting method is to define

$$\chi_0^2 = \sum_{i=1}^{N} \frac{(D_i - T_i)^2}{\sigma_i^2} \qquad \begin{cases} D_i = \text{ data point} \\ T_i = \text{ theory value} \\ \sigma_i = \text{ “expt. error”} \end{cases}$$

and minimize χ_0^2 with respect to the PDF model parameters $\{a_\lambda\}$.

However, the systematic errors imply

$$D_i = T_i(a) + \alpha_i r_{\text{stat},i} + \sum_{k=1}^{K} r_k \beta_{ki}$$

where

α_i = uncorrelated error on D_i

β_{ki} = k^{th} systematic error on D_i

(numbers published by the expt.)

($r_{\text{stat},i}$ and r_k are random variables with standard deviation 1.)

To take into account the systematic errors we define

$$\chi'^2(a_\lambda, r_k) = \sum_{i=1}^{N} \frac{(D_i - \sum_k r_k \beta_{ki} - T_i)^2}{\alpha_i^2} + \sum_k r_k^2,$$

and minimize with respect to both $\{r_k\}$ (≡ the systematic shifts) and $\{a_\lambda\}$ (≡ the PDF model parameters).

Because we use a quadratic penalty term r_k^2, the minimization with respect to $\{r_k\}$ can be done analytically (for arbitrary $\{a_\lambda\}$). Then the minimization w. r. t. $\{a_\lambda\}$ is done numerically.

Comparison of CTEQ6 and CTEQ5



The curves show CTEQ6M and CTEQ5M1 PDF's at $Q=2$ GeV.

(unlabeled : $\bar{u}$ and $s = \bar{s}$; MS-bar scheme)

►Quarks have not changed much.

►Gluon is noticeably different.

The Gluon Distribution



CTEQ6M and CTEQ5M1 gluon distributions at Q = 2 and 100 GeV. (a) Small-x region; (b) large-x region.

► ***Note the hard gluon distribution in CTEQ6M.***

Comparison to Data

χ^2 values for data sets with published correlated systematic errors

data set	N_e	χ^2_e/N_e
BCDMS p	339	1.114
BCDMS d	251	1.114
H1a	104	0.948
H1b	126	1.024
ZEUS	229	1.147
NMC F2p	201	1.517
NMC F2d/p	123	0.909
DØ jet	90	0.766
CDF jet	33	1.472

There is good qualitative agreement between theory and data, but quantitative tests are necessary to assess the uncertainties.

Consider ZEUS F_2 measurements as an example …



CTEQ6M model and ZEUS data in separate x bins. The data points include the optimal shifts for systematic errors. The error bars are statistical errors only.

ZEUS collaboration: S. Chekanov *et al*, Eur Phys J, **C21** (2001) 443.

The "pull" distribution



Histogram of residuals for the ZEUS data.
The curve is a Gaussian of width 1.

(a) $_i = (D_i^{\text{shifted}} - T_i)/\alpha_i$ where $D_i^{\text{shifted}} = D_i - \sum_{k=1}^{N} \hat{r}_k \beta_{ki}.$

(b) The comparison without the systematic shifts.

The optimal systematic shifts $\hat{r}_k$ have reasonable values …

k	1	2	3	4	5	6	7	8	9	10
$\hat{r}_k$	1.67	−0.67	−1.25	−0.44	−0.00	−1.07	1.28	0.62	−0.40	0.21

Tevatron $p\bar{p} \rightarrow jet$ inclusive cross section





The CTEQ6M fit to the inclusive jet data.

(a) DØ data for 5 rapidity bins (0.0 0.5 1.0 1.5 2.0 3.0);

(b) CDF data for central rapidity ($0.1 < |\eta| < 0.7$).

DØ Collaboration: B. Abbott *et al* ; CDF Collaboration: T. Affolder *et al.*

Closer comparison (data-theory / theory) between CTEQ6M and the jet cross section

DØ jet cross section



CDF jet cross section



The Tevatron inclusive jet cross section implies a hard gluon distribution, i.e., $g(x,Q)$ is large at large x.

(Recall CTEQ4HJ and CTEQ5HJ.)

Quantitative Uncertainties

Related Work

S. I. Alekhin, Eur. Phys. J. **C 10** (1999) 395 [hep-ph/9611213]; hep-ex/0005042.

M. Botje, Eur. Phys. J. **C 14** (2000) 285 [hep-ph/9912439].

V. Barone, C. Pascaud and F. Zomer, Eur. Phys. J. **C 12** (2000) 243 [hep-ph/9907512]; C. Pascaud and F. Zomer, Tech. Note LAL-95-05.

A. D. Martin, R. G. Roberts, W. J. Stirling, R. S. Thorne, Eur.Phys.J. **C14** (2000) 133.

W. T. Giele and S. Keller, Phys. Rev. **D 58** (1998) 094023 [hep-ph/9803393]; W. T. Giele, S. A. Keller and D. A. Kosower, "Parton Distribution Function Uncertainties," [hep-ph/0104052].

J. Pumplin, D. R. Stump and W. K. Tung, Phys. Rev. **D 65** (2002) 014011 [hep-ph/0008191].

J. Pumplin *et al.*, Phys. Rev. **D 65** (2002) 014013 [hep-ph/0101032].

D. R. Stump *et al.*, Phys. Rev. **D 65** (2002) 014012 [hep-ph/0101051].

Quantitative Uncertainties

The simplest idea is to plot χ^2_{global} versus the observable of interest, X.



However, because of systematic errors—both experimental and theoretical—normal confidence levels do not apply. Indeed, we conclude that a *large tolerance* (χ^2~100 for 1800 data points) is realistic.

Important computational developments

- Lagrange Multiplier Method (constrained fitting)
- Hessian Matrix Method

  → a complete set of allowed variations using the eigenvector basis

Uncertainty range of the gluon distribution function (at $Q = 2$ GeV).



solid : CTEQ6M
dashed : CTEQ5M1
dotted : MRST 2001
Uncertainty band : envelope of allowed variations

Uncertainties of the parton-parton luminosity functions at the LHC





$$\mathcal{L}(\hat{s}) = \sum_{i,j} C_{ij} \int f_i(x_1) f_j(x_2) \delta(\hat{s} - x_1 x_2 s) dx_1 dx_2$$

provides simple estimates of PDF uncertainties at the LHC.

Example. $\alpha_S(M_Z)$ from the global analysis

Construct *global fits* over a range of α_S. From these alternative models, each data set gives a best value of α_S (min. χ^2) and an "allowed range" of α_S (χ^2 1).



Particle Data Group 0.117 ± 0.002 (shaded strip)

The fluctuations are larger than expected for normal statistics. The vertical lines have $\chi^2_{\text{global}} = 100$,

$$\alpha_S(M_Z) = 0.1165 \pm 0.0065.$$

Determination of α_S from the CTEQ global fit

Plot χ^2 versus α_S for the individual data sets.