


Abstract 376




# Isolated Prompt Photon Production in Photon-Photon Collisions at $\sqrt{s_{ee}}=183-209$ GeV

---

T. Schörner-Sadenius  
on behalf of the OPAL Collaboration


ICHEP 2002, Amsterdam  
23-31 July 2002


'Prompt'  $\equiv$  not coming from particle decays/fragmentation, but emitted from the hard underlying scattering.



# Prompt Photons in $\gamma\gamma \rightarrow \gamma + X$

- Allow to study photon's hadronic structure and test QCD predictions:
  - Prompt  $\gamma$  appear in **resolved processes** (in LO; in NLO also in direct events).
  - Small NLO corrections** / scale dependency.
  - '**Cleaner**' than **dijets** (hadrons,  $D^*$ )  
→ less uncertainty from hadronisation (but prompt  $\gamma$  have smaller Xsection  $\sim O(\alpha_{EM})$ ).
- Might further be used for
  - studies of the underlying **parton picture** or
  - background studies** in LHC Higgs searches.







# Theoretical Predictions


- Fixed-order NLO QCD calculations (Fontannaz et al).
- PYTHIA 6.130 for signal  $\gamma\gamma \rightarrow \gamma + X$ 
  - Distinct samples for single/double resolved for three energy points.
  - Sas-1D (LO)  $\gamma$  PDF.
- Background with fake / real  $\gamma$ :
  - PHOJET 1.10 and HERWIG 5.9 for  $\gamma\gamma$  background without prompt  $\gamma$ .
  - Also Vermaseren 1.0, PYTHIA 5.7, KORALZ, grc4f.

## Single-resolved




## Double-resolved







# LEP and the OPAL Detector

### OPAL Online Data-Taking Statistics




Luminosity used: 639  $\text{pb}^{-1}$






# Event/ $\gamma$ Selection ( $\sqrt{s_{ee}}=183-209$ GeV)

- Require  $\geq 3$  tracks. Reject events with scattered leptons above  $\theta=33$  mrad.
- Reject  $e+e^-$  annihilations:
  - $5 \text{ GeV} < W_{\text{vis}} < 0.3\sqrt{s_{ee}}$
  - $E_{\text{cal}} < 0.25\sqrt{s_{ee}}$ .
- Select isolated photons:
  - EM clusters with  $|\eta| < 1$  and  $p_{\text{T}} > 3 \text{ GeV}$ .
  - Photon and hadronic final state balanced in  $\phi$ .





# Photon Selection ctd'

- Isolation criterion ( $\delta < R$ ):

$$\sum_i E_{Ti} \cdot \Theta(\delta - R_{i\gamma}) \leq 0.2 \cdot E_{T\gamma} (1 - \cos\delta) / (1 - \cos R)$$

- Background: antineutrons,  $\eta/\pi^0$  decays  $\rightarrow$  shape analysis:



$$f_{\max} = E_{\max} / E_{\gamma}$$


$$\sigma_c = (\sum_i E_i R_{i\gamma}) / E_{\gamma}$$

$$g(f, \sigma) = ag^{\gamma} + bg^{\pi} + (1-a-b) \cdot (cg^n + (1-c)g^{\eta})$$

- Fit:  $a = 0.85 \pm 0.08$  (stat)  
 $b = 0.11 \pm 0.08$  (stat)

OPAL preliminary





# Single/Double Resolved Ratio $r_{s/d}$



- For the  $\gamma$  +jet sample:


$$x_{\gamma}^{\pm} = \frac{(E_{\gamma} \pm p_{z\gamma}) + (E_{\text{jet}} \pm p_{z\text{jet}})}{\sum_{\text{hadrons}, \gamma} (E \pm p_z)}$$

using a cone jet finder with  $R=1$ ,  $E_T > 3 \text{ GeV}$ .

- Fit  $x_{\gamma}^{\text{min}}$  distribution to data  $\rightarrow r_{s/d} = 0.59 \pm 0.12$
- For the complete data set use also  $x_T = 2p_T^{\gamma}/W_{\text{vis}}$   
 $\rightarrow r_{s/d} = 0.41 \pm 0.08$ ,
- Average  $\rightarrow r_{s/d} \sim 0.5$ .

OPAL preliminary





# The Total Cross-Section

For isolated prompt photons ( $p_T > 3 \text{ GeV}$ ,  $|\eta^\gamma| < 1$ ):


$$\sigma_{\text{tot}} = r \cdot N_{\text{prompt}} / \varepsilon_{\text{single}} L + (1-r) \cdot N_{\text{prompt}} / \varepsilon_{\text{double}} L$$

- $N_{\text{prompt}}$ : 92.8, background subtracted (FSR,  $\pi$ ,  $\eta$ ,  $n$ ).
- $r_{s/d}$ : single / double resolved ratio.
- $\varepsilon$ : Taken from PYTHIA MC (0.51 / 0.61).



$$\sigma_{\text{tot}} = 0.26 \pm 0.04 \text{ (stat.)} \pm 0.03 \text{ (syst.) pb}$$

- Tristan:  $\sigma(p_T > 2 \text{ GeV}) = 1.72 \pm 0.67 \text{ pb}$  at  $\sqrt{s_{ee}} = 58 \text{ GeV}$








# Comparisons with PYTHIA




→ Shape of distributions well described, but **normalisation off** (1.85; in case of TOPAZ ~3, MC also low for ZEUS)



# Comparisons with NLO Calculations



- Shape and normalisation correct (Fontannaz et al!)



# Summary

---

- Prompt photons provide means to test QCD and to determine hadronic structure of photon.  
 $\sigma = 0.26 \pm 0.04 \text{ (stat.)} \pm 0.03 \text{ (syst.) pb}$
- PYTHIA badly normalized; good agreement of differential cross-sections with NLO QCD calculations.
- Looking forward to final result:
  - Test of different photon PDFs (GRV, LAC1).
  - Investigate including FSR in signal  $\rightarrow$  better NLO description?